

Coherent Beam-Beam Tune Shift of Unsymmetrical Beam-Beam Interactions with Large Beam-Beam Parameter

Lihui Jin and Jicong Shi

Department of Physics & Astronomy, The University of Kansas, Lawrence, KS 66045

Georg H. Hoffstaetter

Department of Physics, Cornell University, Ithaca, NY 14850

Abstract

Coherent beam-beam tune shift of unsymmetrical beam-beam interactions was studied experimentally and numerically in HERA where the lepton beam has a very large beam-beam parameter (up to $\xi_y = 0.272$). Unlike the symmetrical case of beam-beam interactions, the ratio of the coherent and incoherent beam-beam tune shift in this unsymmetrical case of beam-beam interactions was found to decrease monotonically with increase of the beam-beam parameter. The results of self-consistent beam-beam simulation, the linearized Vlasov equation, and the rigid-beam model were compared with the experimental measurement. It was found that the coherent beam-beam tune shifts measured in the experiment and calculated in the simulation agree remarkably well but they are much smaller than those calculated by the linearized Vlasov equation with the single-mode approximation or the rigid-beam model. The study indicated that the single-mode approximation in the linearization of Vlasov equation is not valid in the case of unsymmetrical beam-beam interactions. The rigid-beam model is valid only with a small beam-beam parameter in the case of unsymmetrical beam-beam interactions.

PACS: 29.27.Bd, 29.20.-c, 29.27.-a, 05.45.-a

Corresponding author:

Jack Shi, Department of Physics & Astronomy, University of Kansas, Lawrence, KS 66045

Phone: 785-864-5273; Fax: 785-864-5262; Email: jshi@ku.edu

I. INTRODUCTION

To achieve a substantial increase of luminosity in a storage-ring collider, limited options include increase of bunch currents, reduction of beam sizes at interaction points (IPs), and increase of the number of colliding bunches. The first two measures unavoidably increase head-on beam-beam forces which could lead to collective (coherent) beam-beam instabilities [1, 2]. Understanding of coherent beam-beam effects especially in the nonlinear regime is therefore of primary importance for achieving high luminosity in a storage-ring collider with high-intensity beams.

To study the coherent beam-beam effect, one important quantity that can be measured experimentally is the coherent beam-beam tune shift. Without beam-beam interactions and without considering nonlinearities in the lattice, the two counter-rotating beams oscillate transversely with frequencies that correspond to lattice tunes (betatron tunes without collision) if they deviate from close orbits. With beam-beam interactions, the particle distributions of the beams are perturbed and evolve with time according to the Vlasov equation [1]. The dynamics of the beams could therefore be complicated by multi-mode oscillations of the beam distributions. If considering only the stable oscillation of beam centroids (coherent dipole oscillation), the frequency spectrum of the beam-centroid oscillation has two primary frequencies for each degree of freedom of the transverse motion. These primary frequencies correspond to the tunes measured during collision. The differences between these measured collision tunes and the lattice tunes are the coherent beam-beam tune shifts [1, 3, 4]. Over decades, many studies have been conducted on the relationship between the coherent beam-beam tune shift and the beam-beam parameter that measures the strength of the beam-beam interaction [3-13]. Two theoretical models, the linearized Vlasov equation [1] and the rigid-beam model [5], have been studied extensively for cases of weak beam-beam perturbation in which the beam-beam parameter is relatively small. When the two beams have the same or very close lattice tunes, the calculation of the coherent beam-beam tune shift based on the linearized Vlasov equation with single-mode approximation agrees with beam measurements and computer simulations [3, 4, 6, 7]. The rigid-beam model is inconsistent with the linearized Vlasov equation

and was therefore proven to be wrong by beam measurements in this case [6, 7]. When the two beams have very different lattice tunes, on the other hand, the calculation based on the rigid-beam model provides a good agreement with beam measurements [5, 8, 14]. In both of these models, the equilibrium beam distributions were assumed to be Gaussian distributions for easing the calculations. In the case of weak beam-beam perturbation, this assumption is fairly good as the beams were observed to stay close to a Gaussian.

The situation of strong beam-beam perturbations with relatively large beam-beam parameter is much more complicated and less understood. When the beam-beam parameter exceeds a threshold, the beam-beam interaction could induce a chaotic coherent beam-beam instability. After the onset of the instability, the closed orbits could become unstable for the beam centroids and two beams could develop a spontaneous unstable coherent oscillation [2, 15]. When the beam-beam parameter is below the beam-beam threshold, the coherent beam oscillation is stable. It is, however, not clear whether the linearized Vlasov equation or the rigid-beam model are still valid in the regime of strong beam-beam perturbation. As many efforts are being made to further increase the beam-beam parameter in upgrades of current and developments of future storage-ring colliders, an understanding of the coherent beam-beam tune shift in this regime not only is necessary for the interpretation of the tune measurement during operation of colliders with high-intensity beams but also could shed light on the onset of the chaotic coherent beam-beam stability.

To explore the beam-beam effect with a large beam-beam parameter a beam experiment, the HERA 2000 beam study, was performed on HERA [Hadron Electron Ring Accelerator at DESY (Deutsches Elektronen-Synchrotron), Hamburg, Germany] in which a 920 GeV proton (p) beam and a 27.5 GeV positron (e^+) beam collided at two IPs, H1 and ZEUS [16]. The beam-beam interaction in HERA is a typical case of unsymmetrical beam-beam interaction as the two beams have very different lattice tunes and beam-beam parameters (strong p beam and weak e^+ beam). In the experiment, the vertical beam-beam parameter of the e^+ beam was varied from 0.068 to 0.272 by changing the vertical beta-function of the e^+ beam at two IPs. The emittance of the e^+

beam and the luminosity were measured as functions of the beam-beam parameter. One important phenomenon observed in this experiment is that the measured coherent beam-beam tune shifts of the e^+ beam are much smaller than those calculated from the rigid-beam model. This is the first experimental evidence indicating that the traditional models of the coherent beam-beam tune shift are no longer valid in the situation of strong beam-beam perturbations.

To have a better understanding of the experimental data, we reconstructed the HERA beam experiment with a self-consistent beam-beam simulation. Remarkable agreement between the experiment and the simulation was observed on emittance growth and luminosity reduction. More significantly, the computer simulation confirmed the experimental result of very small coherent beam-beam tune shifts in this case of a very large beam-beam parameter. To examine the validity of the theoretical models for the coherent beam-beam tune shift, the linearized Vlasov equation and the rigid-beam model were solved for the HERA experiment. Since the distribution of the e^+ beam significantly deviated from the Gaussian due to the strong beam-beam interaction, the solutions of the linearized Vlasov equation and the rigid-beam model were calculated with the beam distributions obtained from the beam-beam simulation instead of assuming Gaussian distributions. It was found that for the unsymmetrical beam-beam interaction with a large beam-beam parameter, the result of the rigid-beam model is inconsistent with the beam experiment and beam simulation even though a more accurate beam distribution was used in the calculation. The coherent beam-beam tune shifts calculated from the linearized Vlasov equation with the single-mode approximation were also found to be significantly different from the result of the beam experiment and beam simulation no matter whether the beam-beam parameter is large or small. As the linearization of the Vlasov equation is expected to be valid for at least a small beam-beam parameter, this discrepancy suggests that the single-mode approximation used for solving the linearized Vlasov equation may not be valid in the case of unsymmetrical beam-beam interactions. Unfortunately, without the single-mode approximation the linearized Vlasov equation for the problem of beam-beam interactions is currently unsolvable computationally due to the unsolved degeneracy problem of a matrix with

mode coupling.

This paper is organized as follows. Section II summarizes the results of the HERA 2000 beam experiment. In Section III, the self-consistent beam simulation for the HERA beam experiment is discussed. In Section IV, the coherent beam-beam tune shifts calculated by using the rigid-beam model or the linearized Vlasov equation are compared with the experiment/simulation results. The details of the coherent tune calculation with the theoretical models are presented in Appendix A and B. The characteristics of the coherent beam-beam tune shift in the unsymmetrical case of beam-beam interactions are discussed in Section V. Section VI contains a summary remark.

II. HERA 2000 BEAM EXPERIMENT

In the luminosity upgrade of HERA, the beam-beam parameters of the electron beam have been nearly doubled. To examine any possible luminosity reduction due to beam-beam effects, a series of beam experiments were performed in HERA [14, 16]. In the HERA 2000 beam experiment, the e^+ beam was used to collide with the p beam at the two IPs and the effect of a large beam-beam parameter of the lepton beam was explored by increasing the vertical beta-function ($\beta_{e,y}$) of the e^+ beam at the IPs. The vertical beam-beam parameter ($\xi_{e,y}$) of the e^+ beam is related to $\beta_{e,y}$ by [17]

$$\xi_{e,y} = \frac{r_e N_p}{2\pi\gamma_e} \frac{\beta_{e,y}}{\sigma_{p,y}(\sigma_{p,x} + \sigma_{p,y})} \quad (1)$$

where r_e and γ_e are the classic radius and Lorentz factor of positron, respectively, and N_p is the number of protons per bunch. The horizontal and vertical size of the p beam at the IPs are given by $\sigma_{p,x}$ and $\sigma_{p,y}$. In the experiment, the p beam current (I_p) was fixed. Since the beam-beam parameter of the p beam is very small, there was little change in the p beam size as it was observed during the experiment. The vertical beam-beam parameter $\xi_{e,y}$ is therefore linearly proportional to $\beta_{e,y}$ in this case. During the experiment, after the proton current was filled, $\xi_{e,y}$ was increased from 0.068 to 0.272 as $\beta_{e,y}$ was changed from 1.0 to 4.0 m while other lattice parameters were kept as constants. It should be noted that with two IPs in the HERA experiment a beam-beam parameter of 0.272

is among the highest ever achieved in storage-ring colliders. Table 1 lists the e^+ beam current (I_e) and the beam-beam parameters of the e^+ and p beam at $\beta_{e,y}$ where the measurement was performed. For the p beam, $I_p = 90$ mA, the beta-function at the IPs were $(\beta_{p,x}, \beta_{p,y}) = (7.0, 0.5)$ m; the lattice tunes were $(\nu_x, \nu_y) = (31.291, 32.297)$; and the emittance without collision was $(\epsilon_{0x}, \epsilon_{0y}) = (3.82, 3.18)$ nm. For the e^+ beam, $\beta_{e,x} = 2.5$ m, $(\nu_x, \nu_y) = (52.169, 52.246)$, and $(\epsilon_{0x}, \epsilon_{0y}) = (32.0, 1.28)$ nm. Other beam parameters can be found in [16]. The e^+ and p beam sizes were not matched during the collision. In the experiment, the emittance of the e^+ beam and the luminosity were measured as functions of $\beta_{e,y}$ at both the IPs. In Figs. 1 and 2, the measured emittance and the specific luminosity were plotted, with discrete points, as functions of $\xi_{e,y}$. For each $\xi_{e,y}$ where the measurement was performed, two data points correspond to the measurement at the two IPs, respectively. The specific luminosity is defined as $\mathcal{L}_s = N_{col}\mathcal{L}/(I_e I_p)$ where N_{col} and $\mathcal{L}$ are the number of colliding bunches and the luminosity, respectively [14]. As shown in Fig. 1, the vertical emittance growth of the e^+ beam increases monotonically and smoothly with the increase of $\xi_{e,y}$. This is the characteristics of the incoherent beam-beam effect in contrast to the coherent beam-beam effect of which the emittance growth as a function of the beam-beam parameter could experience certain jumps (phase transitions) due to the onset of the coherent beam-beam instability [2]. To confirm that the luminosity reduction in Fig. 2 is indeed due to the emittance blowup, the luminosity calculated with the measured emittance by using the standard formula [16] is also plotted in the Fig. 2. The agreement between the measured luminosity and the calculated luminosity in Fig. 2 shows a consistency in the emittance and luminosity measurement.

In the experiment, the collision tunes of the e^+ beam were also measured at $\beta_{e,y} = 4.0$ m as $\nu_x = 52.160$ and $\nu_y = 52.233$. The coherent beam-beam tune shift of the e^+ beam is therefore only $\Delta\nu_x = 0.009$ and $\Delta\nu_y = 0.013$ while from the rigid-beam model $\Delta\nu_x = 0.016$ and $\Delta\nu_y = 0.042$ if both the beams are Gaussian [14]. The measured coherent beam-beam tune shifts in this case are inconsistent with the traditional understanding of beam coherent oscillation. Moreover, in the symmetrical case of beam-beam interactions, the ratio of the coherent and incoherent beam-beam tune shift has a

value approximately ranging from 1.2 for round beams to 1.3 for flat (one-dimensional) beams [4]. Since the beam-beam parameters of the p beam is very small, the distribution of the p beam in the HERA experiment is still very close to a Gaussian. The incoherent beam-beam tune shifts that are the tune shifts of a single particle with an infinitesimal betatron oscillation are $2\xi_{e,x} = 0.082$ and $2\xi_{e,y} = 0.544$ for the e^+ beam in the horizontal and vertical plane, respectively. The ratios of coherent and incoherent beam-beam tune shifts of the e^+ beam are $\Delta\nu_{e,x}/(2\xi_{e,x}) = 0.11$ and $\Delta\nu_{e,y}/(2\xi_{e,y}) = 0.024$ at $\beta_{e,y} = 4.0$ m. The coherent beam-beam tune shifts in this case of unsymmetrical beam-beam interactions with a large beam-beam parameter are therefore extremely small as compared with the incoherent beam-beam tune shifts or compared with the symmetrical case of beam-beam interactions.

III. RECONSTRUCTION OF HERA BEAM EXPERIMENT WITH NUMERICAL SIMULATION

To have a better understanding of the measured data in the HERA experiment, we have reconstructed the experiment with a self-consistent beam-beam simulation. This numerical study also served as a detailed benchmark of our beam-beam simulation code with the experimental measurement. In the simulation, the linear HERA lattice with the two IPs was used. The code used in this study is an expanded version of [2] that is currently capable of studying beam-beam effects of proton or lepton beams with any aspect ratio (ratio between vertical and horizontal beam size). The two colliding beams were represented by a million macro-particles with given initial Gaussian distributions in transverse phase space. Without beam-beam interactions, the initial beam distribution used in the simulation matches exactly with the lattice. Beam-beam interaction at each the IP was represented by a kick in transverse phase space and the kick was calculated by using the particle-in-cell method as described in Ref. [2]. Since the beams in the HERA experiment were flat, a uniform mesh extending to $\pm 20\sigma$ in the configuration space with a grid constant of 0.2σ was necessary in this case. Tracking of particle motion was conducted in four-dimensional trans-

verse phase space without synchrotron oscillations and momentum deviations. For lepton beams, the quantum excitation and synchrotron damping were treated as kicks in each turn during the tracking. The horizontal kick is [18]

$$\begin{aligned}\Delta x &= e^{-1/(2\tau_x)}x + \left[\left(1 - e^{-1/(2\tau_x)}\right)\epsilon_x\right]^{1/2}w_1 \\ \Delta p_x &= e^{-1/(2\tau_x)}p_x + \left[\left(1 - e^{-1/(2\tau_x)}\right)\epsilon_x\right]^{1/2}w_2\end{aligned}\tag{2}$$

where x and p_x are the normalized horizontal coordinate and its conjugate momentum, ϵ_x is the horizontal emittance, and w_1 and w_2 are random numbers with a Gaussian distribution that is centered at zero and has unit standard deviation. The damping time in horizontal and vertical direction, τ_x and τ_y , have the unit of turns. For HERA, $\tau_x = 436$ and $\tau_y = 600$, respectively. The vertical kick has a similar formula.

With the beam-beam simulation, the emittance growth of the e^+ beam and the specific luminosity were calculated as functions of $\xi_{e,y}$ for the HERA experiment and the results were plotted in Figs. 1 and 2 as solid lines. Both the emittance and the luminosity plot show a remarkable agreement between the experiment and the simulation. Figure 3 plots the evolution of the vertical emittance of the e^+ beam at different $\beta_{e,y}$. In all these cases, after a quick emittance blowup, the beam emittance is re-stabilized and, consequently, an equilibrium (or quasi-equilibrium) state of the e^+ beam was reached. During the emittance blowup, the particles in the beam core escape quickly to the beam tails. Without the onset of the coherent beam-beam instability, the particles in the beam tails are stable for the beam-beam interaction. This re-stabilization of the beam emittance is therefore due to a depopulation of the beam core and is consistent with the experimental observation that the beam lifetime and operation conditions were good during the experiment even in the case of $\xi_{e,y} = 0.272$ [16].

In the beam simulation, the coherent beam-beam tune shift was also calculated. Fig. 4 is the calculated power spectrum of the coherent oscillation of the e^+ beam at $\beta_{e,y} = 4.0$ m. Due to the quantum fluctuation, the e^+ beam has always a very small oscillation which is enough for

the calculation of the coherent frequency if the numerical noise is small. Since a large number of particles were used for each of the beams, in this simulation the numerical noise was very low and we were able to calculate the coherent frequency without applying off-center kicks on the beams. The spectrum was calculated by the fast Fourier transformation (FFT) of the beam centroid motion from the 5000th to 9000th turn. Since at the 5000th turn the beam has already reached its quasi-equilibrium, the transient state of the beam was thrown out during the tune calculation (see Fig. 3). As shown in Fig. 4 the power spectrum peaks at 0.1605 and 0.2331 in the horizontal and vertical planes, respectively, which corresponds to $\nu_x = 52.161$ and $\nu_y = 52.233$ for the beam coherent tunes during collision. This simulation result agrees excellently with the experimental measurement of $\nu_x = 52.160$ and $\nu_y = 52.233$. The beam-beam simulation therefore confirmed the coherent beam-beam tune shift measured in the HERA beam experiment.

To understand the very small coherent beam-beam tune shift in this case of very large beam-beam parameter, the dynamics of the beam particle distributions was studied during the beam simulation. As shown in Fig. 5, the distribution of the e^+ beam deviates from a Gaussian distribution with a significant drop at the beam core and a growth of the beam tails. Compared with the distribution of the e^+ beam, a Gaussian beam that has the same emittance of the e^+ beam has more particles in the core. The real coherent beam-beam tune shift measured in the experiment and calculated with the beam simulation is therefore smaller than that calculated from the rigid-beam model with Gaussian beams. Moreover, the assumption of Gaussian beams in all the theoretical models of the coherent beam-beam oscillation neglects the effect of the ϕ -dependence of the beam equilibrium distributions, where ϕ is the angle of the action-angle variable. In the HERA beam experiment, the equilibrium (quasi-equilibrium) distribution of the e^+ beam in fact has a strong ϕ -dependence as shown in Fig. 6. For a comparison, the initial distribution used in the simulation was also plotted in the Figure and shows no ϕ -dependence as it should be. In order to compare the experimental/simulation result with calculations of theoretical models, the theoretical models need to be modified to include the ϕ -dependence of non-Gaussian equilibrium distributions.

IV. VALIDITY OF THEORETICAL MODELS FOR BEAM COHERENT OSCILLATION

Two theoretical models, the rigid-beam model and the linearized Vlasov equation with the single-mode approximation, were examined with the HERA experiment. For the rigid-beam model, the coherent tunes were calculated with two different methods: assuming a Gaussian distribution as the equilibrium distribution of the e^+ beam [Eqs. (A10) and (A14) in Appendix A.1] or using the non-Gaussian quasi-equilibrium distribution of the e^+ beam obtained in the beam-beam simulation (Appendix A.2). In the case of the Gaussian distribution, the beam sizes used in Eq. (A14) are of the experimental measurement or the beam-beam simulation. The small differences in the beam sizes measured in the experiment or calculated from the simulation (see Fig. 1) made little difference in Eq. (A14). For the linearized Vlasov equation, the horizontal coherent tunes were obtained by solving the initial-value problem of the linearized Vlasov equation with the single-mode approximation in the horizontal plane. The details of the calculations are in Appendix A and B.

Tables II and III list the ratio of the coherent and incoherent beam-beam tune shifts of the e^+ beam, $\Delta\nu_{e,x}/(2\xi_{e,x})$ and $\Delta\nu_{e,y}/(2\xi_{e,y})$, calculated in the beam simulation or calculated with the theoretical models for the cases of $\beta_{e,y} = 1.0$ and 4.0 m, where $\Delta\nu_{e,x}$ and $\Delta\nu_{e,y}$ are the horizontal and vertical coherent beam-beam tune shift of the e^+ beam. The experimental measurement at $\beta_{e,y} = 4.0$ m is also included in Table II. The significant discrepancy between the results of the models and the results of the experiment/simulation shows that the theoretical models are inconsistent with the experiment and simulation. Note that the rigid-beam model with the beam distribution from the simulation did a little better than that with Gaussian distribution. To further examine the failure of the theoretical models, a beam-beam simulation was conducted for the case of $\beta_{e,y} = 1.0$ m but with only one tenth of the p -bunch current used in the experiment, i.e. $\xi_{e,y} = 0.0068$. The result is listed in Table IV and shows that the rigid-beam model with either the Gaussian distribution or the distribution obtained from the simulation are in a good agreement with the beam simulation. In fact, the quasi-equilibrium distribution of the e^+ beam in this case

is very close to a Gaussian. This re-confirms many previous studies that the rigid-beam model is correct for unsymmetrical beam-beam interactions with relatively small beam-beam parameter.

In the case of the linearized Vlasov equation, as shown in Tables II-IV, the calculation yielded wrong results no matter how small the beam-beam parameter is. In the calculation with the Vlasov equation, several approximations were employed. Among them, the linearization, the one-dimensional beam, and the single-mode approximation are the three major approximations that cannot be directly justified by the experimental observations (see Appendix B). The linearization of the Vlasov equation should not play the leading role in its failure for the unsymmetrical beam-beam interaction since the linearization should be valid for a relatively small beam-beam parameter. To verify the validity of the one-dimensional approximation, we did a series of beam-beam simulations for the case of $\beta_{e,y} = 1.0$ m, $\xi_{e,y} = 0.068$ or 0.0068 , but with a different beam aspect ratio ($\sigma_{y,e}/\sigma_{x,e}$) ranging from 0.063 to 0.120 . In all these cases, the horizontal coherent beam-beam tune shift was found to be very similar to that of the HERA experiment. Note that in the HERA experiment and in all the cases in Tables II-IV, $\sigma_{y,e}/\sigma_{x,e} = 0.126$. This indicates that the horizontal coherent beam-beam tune shift is not sensitive to the aspect ratio of the beam. Similar phenomenon has also been observed in the symmetrical case of beam-beam interactions [4]. The large discrepancy in the coherent beam-beam tune shifts calculated with the linearized Vlasov equation as compared to that obtained from the experiment/simulation is apparently not due to the approximation of one-dimensional beam. With the single-mode approximation, on the other hand, the only oscillation mode of the beam distribution that was kept in the calculation is the $m = 1$ mode in Eq. (B6). As shown in Fig. 6, the equilibrium distribution of the e^+ beam [$f_{1,0}(I, \phi)$ in Eq. (B4)] depends strongly on ϕ . In this case, the dominant mode of the equilibrium distribution is the $m = 2$ mode which has a ϕ -dependence of $e^{i2\phi}$. As the $m = 2$ mode could be a dominant intrinsic mode of the system, the single-mode approximation could be inconsistent with the dynamics of the beam. In order to study the effect of mode couplings in the linearized Vlasov equation, we have derived the eigenvalue equation of the linearized Vlasov equation that is similar to Eq. (B13) but includes

high-order modes. After truncating higher-order modes at $m = m_p$, $\mathbf{M}$ in Eq. (B13) becomes a $(2m_p + 1)(2l_p + 2) \times (2m_p + 1)(2l_p + 2)$ matrix where l_p is the number of grids on the mesh of the action space and $\mathbf{M}_1$ and $\mathbf{M}_2$ in Eq. (B14) are no longer diagonal matrices (see Appendix B). We have, however, failed in obtaining a set of orthogonal eigenvectors for the eigenvalue equation of the linearized Vlasov equation because $\mathbf{M}$ is a singular (ill-conditioned) matrix when the mode couplings are included. Similar problem has also been encountered when including the mode coupling in the case of symmetrical beam-beam interactions.

V. COHERENT VERSUS INCOHERENT BEAM-BEAM TUNE SHIFT

In the symmetrical case of beam-beam interactions, the ratio of the coherent and incoherent beam-beam tune shift is approximately a constant of 1.2 for round beams and 1.3 for flat beams [4]. Note that with symmetrical beam-beam interactions, the beam-size growths of the two beams are symmetrical and the beam distributions are usually close to a Gaussian when the beam-beam parameter is below the threshold of the onset of coherent beam-beam instability [2, 15]. The coherent beam-beam tune shift depends linearly on the beam-beam parameter in a fairly large range of beam-beam parameter [4]. On the contrary, in the HERA experiment the ratio of the coherent and incoherent beam-beam tune shift of the e^+ beam decreases monotonically with the increase of the beam-beam parameter as shown in Fig. 7. This different characteristic of the coherent beam-beam tune shift stems mainly from the mismatch in the equilibrium distributions of two unsymmetrical colliding beams. In the HERA experiment, the beam size of the e^+ beam at the IPs is slightly larger than that of the p beam initially without collision (see Table I). When the beam-beam parameter is small such as in the case of $\xi_{e,y} = 0.0068$, the beam-size growth is insignificant and the beams are close to the Gaussians. The ratio of the coherent and the incoherent beam-beam tune shift of the e^+ beam is simply determined by the mismatch in the initial beam sizes without collision as described by Eq. (A14) of the rigid-beam model. When the beam-beam parameter is large, on the other hand, the beam-size growth of the e^+ beam is significant (see Fig.

1) and dominates the beam-size mismatch. Moreover, the particle distribution of the e^+ beam deviates significantly from a Gaussian distribution (see Fig. 5). In this situation, the size of the e^+ beam is much larger than that of the p beam and a large number of e^+ particles escape to the beam tails that is away from the core of the p beam. The coherent beam-beam tune shift of the e^+ beam thus becomes much smaller than that in the case of two matched Gaussian beams. Note that for two matched Gaussian beams, the ratio of the coherent and incoherent beam-beam tune shift is 0.5 for unsymmetrical beam-beam interactions. As the beam-size growth and the resultant beam-size mismatch increase with the beam-beam parameter, the ratio of the coherent and incoherent beam-beam tune shift of the e^+ beam decreases with the beam-beam parameter. In general, the equilibrium states of the Vlasov equation in Eq. (B4) are different for two unsymmetrical colliding beams due to the asymmetry of beam-beam perturbations. The mismatch in the distributions as well as the beam sizes is, in principle, independent of the initial states of the beam distributions and the initial beam-size mismatch as long as the considered equilibrium state of the Vlasov equation in Eq. (B4) is well isolated and is the only one that is close to the initial beam distributions. The beam-size mismatch in the unsymmetrical case of beam-beam interactions is therefore intrinsic and unavoidable especially when the beam-beam parameter is larger. Moreover, this beam-size mismatch increases with the strength of beam-beam perturbations. The functional dependence of the coherent beam-beam tune shift on the beam-beam parameter in Fig. 7 is therefore a general characteristic of unsymmetrical beam-beam interactions.

Figure 7 also plots $\Delta\nu_{e,y}/(2\xi_{e,y})$ as a function of $\xi_{e,y}$ calculated based on the rigid-beam model. It shows that when the beam-beam parameter is large, the rigid-beam model results in a similar $\xi_{e,y}$ -dependence of $\Delta\nu_{e,y}/(2\xi_{e,y})$ even though it overestimates the coherent beam-beam tune shift. Note that in the rigid-beam model, this $\xi_{e,y}$ -dependence is due to the beam-size mismatch as given in Eq. (A14). At a large $\xi_{e,y}$, the beam-size mismatch is dominated by the beam-size growth of the e^+ beam. This suggests that the result of the rigid-beam model in Eq. (A14) provides an approximation of the functional dependence of the coherent beam-beam tune shift to the intrinsic

beam-size mismatch in unsymmetrical beam-beam interactions with large beam-beam parameters.

VI. CONCLUSION

The coherent beam-beam tune shift was studied in the case of unsymmetrical beam-beam interactions where the two beams have very different beam-beam parameters and betatron tunes. The results of a self-consistent beam-beam simulation, the rigid-beam model, and the linearized Vlasov equation were compared with the beam measurement in the HERA 2000 Beam Study. Remarkable agreement was found between the beam simulation and the HERA experiment in a wide range of and, especially, at very large beam-beam parameters of the lepton beam. The rigid-beam model was found to be only correct when the beam-beam parameter is small. The result of the linearized Vlasov equation with the single-mode approximation is inconsistent with the result of the beam experiment/simulation in either cases of large or small beam-beam parameter. The failure of the linearized Vlasov equation could be due to the single-mode approximation used in solving the linearized Vlasov equation. A study of the dynamics of the beam distribution showed that the high-order modes are important to the beam dynamics in this situation. An attempt to include high-order modes in the calculation has, however, not been successful because of the difficulty in finding a set of orthogonal eigenvectors for the linearized Vlasov equation. Recently, efforts have been made to include the angle-dependence of beam distributions in an expansion of the Vlasov equation [19]. More studies are needed for a relevant solution of the linearized Vlasov equation for the unsymmetrical beam-beam interaction. Currently, the numerical simulation is the only reliable approach for an prediction of the coherent beam-beam tune shift in this situation.

One interesting phenomenon observed in this study is the very small coherent beam-beam tune shift in this unsymmetrical case of beam-beam interactions. It was found that the ratio of the coherent and incoherent beam-beam tune shift of the weak lepton beam in HERA decreases from 0.3 to 0.02 as the incoherent beam-beam tune shift increases from 0.01 to 0.54. On the contrary, in the symmetrical case of beam-beam interactions, this ratio maintains approximately a constant

of 1.2 for a round beam or 1.3 for a flat beam in a large range of beam-beam parameter. The reason for this different characteristic of the coherent beam-beam tune shift is the intrinsic beam-size mismatch between two unsymmetrical colliding beams due to the difference in the equilibrium distributions of the two beams. This intrinsic mismatch in the beam distributions due to beam-beam interactions becomes more pronounced as the strength of beam-beam perturbations increases. The ratio of the coherent and incoherent beam-beam tune shift decreases, in general, with the increase of beam-beam parameters in the unsymmetrical case of beam-beam interactions.

APPENDIX A. COHERENT BEAM-BEAM TUNE SHIFT FROM RIGID-BEAM MODEL

The coherent beam-beam tune shift in the rigid-beam model has been studied before for the unsymmetrical beam-beam interaction. Previous studies are, however, limited to the special cases where either the two beams have the same [8] or very different [14] beam-beam parameters. Moreover, all of them assumed that the beams are Gaussian. In the following, a general formula of the rigid-beam model is derived that can be apply to any case of unsymmetrical beam-beam interaction with either Gaussian or non-Gaussian beams.

Let $\rho_i(\vec{r}, \theta)$ be the distribution of beam i in the normalized configuration space where $i = 1$ or 2 , $\vec{r} = (x, y)$ are the normalized coordinates of the transverse space, and θ the azimuthal angle associated with the path length along the closed orbit. The beam centroid in the normalized space can be calculated by $\vec{R}_i = \int \vec{r} \rho_i(\vec{r}, \theta) d\vec{r}$. Considering a linear lattice with one IP, the transverse motion of the beam centroid can be described by

$$\frac{d^2 \vec{R}_i}{d\theta^2} + \mathbf{\Omega}_i \cdot \vec{R}_i = (-1)^{i+1} \lambda_i \vec{F} \left[2\pi \sum_n \delta(\theta - 2\pi n) \right]. \quad (\text{A1})$$

In Eq. (A1), $\mathbf{\Omega}_i$ is a 2×2 diagonal matrix with $(\mathbf{\Omega}_i)_{11} = \nu_{i,x}^2$ and $(\mathbf{\Omega}_i)_{22} = \nu_{i,y}^2$, where $(\nu_{i,x}, \nu_{i,y})$ are the fractional parts of the betatron tunes of the lattice for beam i . The main part of the

beam-beam kick in Eq. (A1) is

$$\vec{F} = \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \rho_1(\vec{r}_1, \theta) \rho_2(\vec{r}_2, \theta) \vec{G}(\beta_{1,x}^{1/2} x_1 - \beta_{2,x}^{1/2} x_2, \beta_{1,y}^{1/2} y_1 - \beta_{2,y}^{1/2} y_2) d\vec{r}_1 d\vec{r}_2 \quad (A2)$$

where $\vec{G}(x, y) = \vec{r}/r^2$ is the Green Function of the beam-beam interaction and $(\beta_{i,x}, \beta_{i,y})$ are the horizontal and vertical beta-function of beam i at the IP, respectively. The strength of the beam-beam kick for the horizontal component of Eq. (A1) is

$$\lambda_i = \lambda_{i,x} = \frac{a_i N_j}{\pi \gamma_i} \nu_{i,x} \beta_{i,x}^{1/2}$$

where $i=1$ or 2 , $j=1$ or 2 , but $i \neq j$; N_j is the number of particles per bunch of beam j ; a_i the classical radius of the particle in beam i ; and γ_i the Lorentz factor of beam i . With the definition of the beam-beam parameters in Eq. (1), this kick strength can be written as

$$\lambda_{i,x} = \frac{2\nu_{i,x} \xi_{i,x}}{\beta_{i,x}^{1/2}} \sigma_{j,x}(\sigma_{j,x} + \sigma_{j,y}) \quad (A3)$$

where $(\xi_{i,x}, \xi_{i,y})$ and $(\sigma_{i,x}, \sigma_{i,y})$ are the beam-beam parameters and the rms beam sizes at the IP of beam i . For the vertical component of Eq. (A1), $\lambda_i = \lambda_{i,y}$ can be easily obtained by exchanging x and y in Eq. (A3).

In the rigid-beam model of the coherent beam-beam oscillation, the shapes of the particle distributions in phase space are assumed to not change with time during the beam oscillation while the centers of the distributions oscillate with the beams' coherent tunes. The distribution during the beam oscillation is thus assumed to be $\rho_i(\vec{r}, \theta) = \rho_{0i}(\vec{r} - \vec{R}_i)$, where $\rho_{0i}(\vec{r})$ is the equilibrium distribution when the beam is centered at the closed orbit. In general, $\vec{F}$ in Eq. (A2) is a function of moments of phase-space variables and the time-dependence of $\vec{F}$ is implicitly through all the moments. (To have a better picture of this, one may consider a moment expansion of beam particle distributions in phase space). With the rigid-beam approximation, $\vec{F}$ depends on the lowest-order moments $\vec{R}_1$ and $\vec{R}_2$ only,

$$\vec{F} = \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \rho_{01}(\vec{r}_1 - \vec{R}_1) \rho_{02}(\vec{r}_2 - \vec{R}_2) \vec{G}(\beta_{1,x}^{1/2} x_1 - \beta_{2,x}^{1/2} x_2, \beta_{1,y}^{1/2} y_1 - \beta_{2,y}^{1/2} y_2) d\vec{r}_1 d\vec{r}_2 \quad (A4)$$

Note that the rigid-beam model may fail in cases where the variation of the distributions is important during the beam coherent oscillation. To find the oscillation frequencies of $\vec{R}_i$, one can average the beam-beam kick in Eq. (A1) over one turn (2π in longitudinal direction) and expand $\vec{F}$ into a Taylor series of $\vec{R}_i$. Keeping only the linear terms of $\vec{R}_i$, Eq. (A1) becomes a coupled four-dimensional harmonic oscillator,

$$\frac{d^2 \vec{R}_i}{d\theta^2} + \mathbf{\Omega}_i \cdot \vec{R}_i = (-1)^{i+1} \lambda_i \left(\mathbf{A}_1 \cdot \vec{R}_1 + \mathbf{A}_2 \cdot \vec{R}_2 \right) \quad (A5)$$

where $i = 1$ or 2 , and $\mathbf{A}_i$ are 2×2 matrices with

$$\mathbf{A}_i = \left. \frac{\partial \vec{F}}{\partial \vec{R}_i} \right|_{\vec{R}_1=0, \vec{R}_2=0}. \quad (A6)$$

If both the beams are mirror symmetric with respect to the horizontal and vertical plane, $\mathbf{A}_i$ are diagonal matrices and the horizontal and vertical coherent oscillation are decoupled. The two eigenfrequencies for the coherent oscillation in the horizontal plane can then be solved as

$$\nu_{\pm} = \frac{1}{\sqrt{2}} \sqrt{\omega_1^2 + \omega_2^2 \pm \sqrt{(\omega_1^2 - \omega_2^2)^2 + 16\nu_{1,x}\nu_{2,x}\delta\omega_1\delta\omega_2}} \quad (A7)$$

where

$$\begin{aligned} \omega_i^2 &= \nu_{i,x}^2 - 2\nu_{i,x}\delta\omega_i \\ \delta\omega_i &= \lambda_{i,x} (\mathbf{A}_i)_{11} / (2\nu_{i,x}) \end{aligned} \quad (A8)$$

for $i = 1$ or 2 . If $\delta\omega_i \ll \nu_{i,x}$ and in Eq. (A7)

$$(\omega_1^2 - \omega_2^2)^2 \gg 16\nu_{1,x}\nu_{2,x}\delta\omega_1\delta\omega_2 \quad (A9)$$

then the coherent frequencies of the two beam are

$$\begin{cases} \nu_+ = \omega_1 = \nu_{1,x} - \delta\omega_1 \\ \nu_- = \omega_2 = \nu_{2,x} - \delta\omega_2 \end{cases} \quad (A10)$$

where ν_+ and ν_- are the horizontal coherent tunes of beam 1 and beam 2, respectively. Note that in the unsymmetrical case of beam-beam interactions, the two eigenfrequencies do not correspond

to the so called 0- (or σ -) and π -modes of symmetrical beam-beam interactions. The condition in Eq. (A9) can be further simplified as

$$|\nu_{1,x} - \nu_{2,x}| \gg \sqrt{\delta\omega_1 \delta\omega_2}. \quad (\text{A11})$$

Therefore, if the difference of the lattice tunes is much larger than the geometric average of the coherent beam-beam tune shifts of the two beams, the coherent beam-beam tune shifts can simply be calculated with Eq. (A10). Note that in the HERA experiment this condition was fulfilled. For the case of strong-weak beam-beam interactions such as $\xi_{1,x} \gg \xi_{2,x}$, one can expand $\nu_{\pm}$ in terms of $\lambda_{2,x}/\lambda_{1,x}$. Keeping only the dominant term in the coherent beam-beam tune shifts yields

$$\begin{cases} \nu_+ = \nu_{1,x} - \delta\omega_1 \\ \nu_- = \nu_{2,x} - \frac{\nu_{1,x}^2 - \nu_{2,x}^2}{\nu_{1,x}^2 - \nu_{2,x}^2 - 2\nu_{1,x}\delta\omega_1} \delta\omega_2 \end{cases} \quad (\text{A12})$$

where ν_+ and ν_- are the coherent frequencies of the weak (beam 1) and strong (beam 2) beam, respectively. In the first equation of Eq. (A12), since the zeroth-order term ($\delta\omega_1$) of $\lambda_{2,x}/\lambda_{1,x}$ exists and dominates the coherent beam-beam tune shift of beam 1, the first- or higher-order terms were neglected. In the second equation of Eq. (A12), on the other hand, the zeroth-order term is zero and the first-order term was thus kept. Note that if the lattice tunes of the two beams are very different, Eq. (A12) is equivalent to Eq. (A10). If the denominator in Eq. (A12), $\nu_{1,x}^2 - \nu_{2,x}^2 - 2\nu_{1,x}\delta\omega_1$, is small, an analysis of the higher order terms shows that the expansion in terms of $\lambda_{2,x}/\lambda_{1,x}$ is no longer accurate and the coherent tunes have to be calculated by using Eq. (A7). The two coherent frequencies in the vertical plane can be easily obtained by exchanging x and y and changing $(\mathbf{A}_i)_{11}$ with $(\mathbf{A}_i)_{22}$ in Eqs. (A7)-(A12). As shown in our study (see Section IV), this approach of the rigid-beam model is quite good in the case of unsymmetrical beam-beam interactions with a small beam-beam parameter.

A.1. Gaussian Beams

In the case that ρ_{0i} are Gaussian distributions, matrix $\mathbf{A}_i$ in Eq. (A6) can be calculated

analytically with Eq. (A4) as

$$(\mathbf{A}_i)_{11} = \frac{\beta_{i,x}^{1/2}}{\Sigma_x(\Sigma_x + \Sigma_y)} \quad (\text{A13})$$

where $\Sigma_x = \sqrt{\sigma_{1,x}^2 + \sigma_{2,x}^2}$ and $\Sigma_y = \sqrt{\sigma_{1,y}^2 + \sigma_{2,y}^2}$. The matrix element of $(\mathbf{A}_i)_{22}$ can be simply obtained by exchanging x and y in Eq. (A13). Substituting Eqs. (A2) and (A13) into Eq. (A8) yields

$$\delta\omega_i = \xi_{i,x} \frac{(\sigma_{j,x} + \sigma_{j,y})\sigma_{j,x}}{(\Sigma_x + \Sigma_y)\Sigma_x} \quad (\text{A14})$$

where $i = 1$ or 2 ; $j = 1$ or 2 ; but $j \neq i$. When $i = 1$, Eq. (A14) gives the coherent beam-beam tune shifts of the weak beam [beam 1 in Eq. (A12)] obtained previously by Hoffstaetter for the strong-weak case of beam-beam interactions [14]. If $(\nu_{1,x}, \nu_{1,y}) = (\nu_{2,x}, \nu_{2,y})$ and $(\sigma_{1,x}, \sigma_{1,y}) = (\sigma_{2,x}, \sigma_{2,y})$, Eq. (A7) is reduced to the formula obtained by Hirata [5]. If $(\xi_{1,x}, \xi_{1,y}) = (\xi_{2,x}, \xi_{2,y})$ and $(\sigma_{1,x}, \sigma_{1,y}) = (\sigma_{2,x}, \sigma_{2,y})$, Eq. (A7) is reduced to the formula obtained by Hofmann [8].

A.2. Non-Gaussian Beams

For non-Gaussian beams, especially the distributions obtained from beam-beam simulations such as that in Figs. 5 and 6, matrices $\mathbf{A}_i$ in Eq. (A6) cannot be obtained analytically but can be calculated numerically by using Eq. (A4). The coherent frequencies can then be calculated with Eq. (A7) or directly from Eq. (A5) if the horizontal and vertical motion are coupled.

APPENDIX B. COHERENT BEAM-BEAM TUNE SHIFT FROM LINEARIZED VLASOV EQUATION

The use of the linearized Vlasov equation has been very successful for the coherent beam-beam tune shift in the case that two beams have the same or very close lattice tunes [4]. In order to find the coherent beam-beam tune shift, one needs to identify the coherent frequencies from the eigenfrequencies of the linearized Vlasov equation. The linearized Vlasov equation, in principle, has infinite numbers of eigenfrequencies associated with infinite numbers of oscillation modes. For real beams, the number of the eigenfrequencies of the beam oscillation in transverse space is twice of the number of particles in a bunch. When two beams have the same lattice tunes,

the coherent frequencies can be easily identified since the eigenfrequencies that correspond to the coherent frequencies are separated from the rest of the eigenfrequencies that form a continuous band (many close eigenfrequencies lines) [4]. Although, it is not very clear mathematically why this separation occurs. The situation becomes more complicated when two beams have very different lattice tunes. In this case, all the eigenfrequencies are in one or two continuous bands and the coherent frequencies cannot be identified by only solving the eigenfrequencies. In order to find the coherent frequencies in the HERA beam experiment, we will instead solve the initial-value problem of the linearized Vlasov equation for the coherent beam oscillation.

Consider only the horizontal motion (very flat beam) in a linear lattice with one IP. In terms of action-angle variable, the Hamiltonian for the betatron motion of beam i ($i = 1$ or 2) can be written as

$$H_i(I, \phi, \theta) = H_{i,0}(I) + U_i(I, \phi, \theta) \left[2\pi \sum_n \delta(\theta - 2\pi n) \right] \quad (B1)$$

where $H_{i,0} = \nu_{i,x}I$ is the Hamiltonian associated with the betatron motion in the linear lattice and U_i is the potential energy for the beam-beam interaction that can be written, for one-dimensional beams, as

$$U_i(I, \phi, \theta) = U_i[f_j] = -2 \frac{\xi_{i,x} \sigma_{j,x}^2}{\beta_{i,x}} \int_0^{2\pi} \int_0^\infty f_j(I', \phi', \theta) \ln \left(\sqrt{2\beta_{i,x}I} \sin \phi - \sqrt{2\beta_{j,x}I'} \sin \phi' \right) dI' d\phi' \quad (B2)$$

where $i = 1$ or 2 , and $j = 1$ or 2 , but $i \neq j$. The action-angle variables are related to the normalized variables by $x = \sqrt{2I} \sin \phi$ and $p = \sqrt{2I} \cos \phi$. $f_i(I, \phi, \theta)$ is the particle distribution of beam i in phase space and satisfies the Vlasov equation. For convenience, we also define a functional $U_i[f_j]$ in Eq. (B2) for the potential integral. In Eq. (B2), $\ln(x - x')$ is the Green function for the potential of beam-beam interaction in one-dimensional space. If only the coherent beam-beam tune shifts are interested, one can get rid of the periodic δ -function in the Hamiltonian in Eq. (B1) by average the beam-beam force over one turn. The Vlasov equation for f_i can then be written as

$$\frac{\partial f_i}{\partial \theta} + \nu_{i,x} \frac{\partial f_i}{\partial \phi} = \{U_i, f_i\} \quad (B3)$$

where $\{ \}$ is the Poisson bracket. Assume that the beams have reached equilibrium distributions $f_{i,0}$ that satisfy

$$\nu_{i,x} \frac{\partial f_{i,0}}{\partial \phi} = \{U_{i,0}, f_{i,0}\} \quad (B4)$$

where $U_{i,0}(I, \phi) = U_i[f_{i,0}]$. Consider that beam i experiences a small perturbation from its equilibrium distribution $\psi_i(I, \phi, \theta) = f_i(I, \phi, \theta) - f_{i,0}(I, \phi)$. The linearized equation for $\psi_i(I, \phi, \theta)$ can be obtained by subtracting Eq.(B4) from Eq. (B3) and neglecting the term $\{U_i[\psi_j], \psi_i\}$ which is higher-order in ψ_i as

$$\frac{\partial \psi_i}{\partial \theta} + \nu_{i,x} \frac{\partial \psi_i}{\partial \phi} = \{U_{i,0}, \psi_i\} + \{V_i, f_{i,0}\}. \quad (B5)$$

where $V_i(I, \phi, \theta) = U_i[\psi_j]$.

To solve Eq. (B5), one can convert it into a system of infinite numbers of coupled ordinary differential equations of modes by using Fourier transformation

$$\psi_i(I, \phi, \theta) = \int_{-\infty}^{\infty} d\nu \sum_{m=-\infty}^{\infty} \psi_{i,m}(I, \nu) e^{i(m\phi - \nu\theta)} \quad (B6)$$

where ν is the oscillation frequency of the beams and m numbers modes. The $m = 1$ mode corresponds to the coherent dipole oscillation. To further simplify the problem, one may use the single-mode approximation in which only the mode with $m = 1$ is kept in the linearized Vlasov equation [4]. It turns out that the use of single-mode approximation is not only a convenience but also a necessity. Without the single-mode approximation, no effective method is available for the general solution of Eq. (B5), except for simplified models as in Ref. [19]. Substituting Eq. (B6) into Eq. (B5), multiplying $e^{-i\phi}$ and integrating ϕ over 2π on the both sides of Eq. (B5), and only keeping the $m = 1$ mode yields

$$\nu \bar{\psi}_i(I, \nu) = \nu_{i,x} \bar{\psi}_i(I, \nu) + Q_i(I) \bar{\psi}_i(I, \nu) + \int_0^{\infty} G_i(I, I') \bar{\psi}_j(I', \nu) dI' \quad (B7)$$

where $\bar{\psi}_i(I, \nu) = \psi_{i,1}(I, \nu)$,

$$Q_i(I) = \frac{1}{2\pi} \int_0^{2\pi} \frac{\partial U_{i,0}}{\partial I} d\phi \quad (B8)$$

and

$$G_i(I, I') = \frac{\xi_{i,x} \sigma_{j,x}^2}{\pi \sqrt{2\beta_{i,x} I}} \int_0^{2\pi} \frac{e^{-i\phi} \sin \phi'}{\sqrt{\beta_{i,x} I} \sin \phi - \sqrt{\beta_{j,x} I'} \sin \phi'} \left(2I \cos \phi \frac{\partial f_{i,0}}{\partial I} - \sin \phi \frac{\partial f_{i,0}}{\partial \phi} \right) d\phi' d\phi \quad (B9)$$

If the equilibrium distributions are independent of ϕ such as for Gaussian beams, the imaginary term of $G_i(I, I')$ is zero. Otherwise, this imaginary term contributes a damping to the linearized Vlasov equation when $\bar{\psi}_i$ is stable or an excitation when $\bar{\psi}_i$ is unstable. If the equilibrium distributions $f_{i,0}$ are Gaussian, with a similar algebraic treatment in Ref. [4], the integrals in Eqs. (B8) and (B9) can be calculated analytically as

$$Q_i(I) = -\frac{\xi_{i,x} \sigma_{j,x}^2}{\beta_{i,x} I} \left(1 - e^{-\beta_{i,x} I / \sigma_{j,x}^2} \right) \quad (B10)$$

$$G_i(I, I') = \xi_{i,x} r_{ij} e^{-(z_i + z'_j)/2} \left[\frac{\min(z_i, r_{ij} z'_j)}{\max(z_i, r_{ij} z'_j)} \right]^{\frac{1}{2}} \quad (B11)$$

where $i = 1$ or 2 ; $j = 1$ or 2 ; but $i \neq j$. $z_i = \beta_{i,x} I / \sigma_{i,x}^2$, $z'_j = \beta_{j,x} I' / \sigma_{j,x}^2$, and $r_{ij} = \sigma_{j,x}^2 / \sigma_{i,x}^2$.

B.1. Eigenfrequencies and Eigenvectors of Linearized Vlasov Equation

To further proceed with Eq. (B7), one may discretize the action space (I) into a mesh and solve the equation on the grids [3, 4]. Let $I = l\Delta I$ where ΔI is the grid size; $l = 0, 1, 2, \dots, l_p$; and $l_p \Delta I$ is the size of the mesh. Since the distributions decay to zero quickly as I increases, a mesh that covers several $\sigma_{i,x}$ is good enough for a calculation of the coherent frequency. In order to have an accurate frequency for the lattice tune in the eigenfrequencies of Eq. (B7), however, the mesh has to be large enough so that the incoherent beam-beam tune shift at $I = l_p \Delta I$ is negligible. In this study, we therefore used $l_p \Delta I = 160 \epsilon_{i,x}$ and $\Delta I = 0.05 \epsilon_{i,x}$, where $\epsilon_{i,x}$ is the normalized emittance of beam i . Let $\bar{\psi}_i(l\Delta I, \nu) = \bar{\psi}_{il}(\nu)$. Eq. (B7) can then be converted into a system of linear algebraic equations on the mesh,

$$\nu \bar{\psi}_{il} = \nu_{i,x} \bar{\psi}_{il} + Q_i(l\Delta I) \bar{\psi}_{il} + \Delta I \sum_{k=0}^{l_p} G_i(l\Delta I, k\Delta I) \bar{\psi}_{jk}. \quad (B12)$$

that leads to an eigenvalue problem

$$\mathbf{M}\vec{V} = \nu \vec{V} \quad (B13)$$

where

$$\vec{V} = \left(\bar{\psi}_{10}, \bar{\psi}_{11}, \dots, \bar{\psi}_{1l_p}, \bar{\psi}_{20}, \bar{\psi}_{21}, \dots, \bar{\psi}_{2l_p} \right)^T.$$

$\mathbf{M}$ is a $2(l_p + 1) \times 2(l_p + 1)$ matrix

$$\mathbf{M} = \begin{pmatrix} \mathbf{M}_1 & \mathbf{O}_1 \\ \mathbf{O}_2 & \mathbf{M}_2 \end{pmatrix} \quad (B14)$$

where $\mathbf{M}_1$, $\mathbf{M}_2$, $\mathbf{O}_1$, and $\mathbf{O}_2$ are $(l_p + 1) \times (l_p + 1)$ matrices. Because of the single-mode approximation, $\mathbf{M}_1$ and $\mathbf{M}_2$ are diagonal matrices with the diagonal elements

$$(\mathbf{M}_i)_{kk} = \nu_{i,x} - Q_i((k - 1)\Delta I) \quad (B15)$$

and the elements of $\mathbf{O}_i$ are

$$(\mathbf{O}_i)_{kl} = G_i((k - 1)\Delta I, (l - 1)\Delta I) \quad (B16)$$

where $k = 1, \dots, (l_p + 1)$; $l = 1, \dots, (l_p + 1)$; and $i = 1$ or 2 . If the equilibrium distributions are Gaussian, all these matrix elements in Eqs. (B15) and (B16) can be calculated analytically by using Eqs. (B10) and (B11). In the case of the HERA experiment, the equilibrium distribution of the p beam is still very close to a Gaussian but the e^+ beam is no longer a Gaussian beam (see Figs. 5 and 6). Let beam 1 and 2 be the e^+ and p beam, respectively. $\mathbf{M}_1$ and $\mathbf{O}_2$ can then be obtained analytically. The matrix elements of $\mathbf{M}_2$ and $\mathbf{O}_1$, on the other hand, have to be calculated numerically by using Eqs. (B8) and (B9) with the quasi-equilibrium distribution of the e^+ beam obtained from the beam simulation.

With the eigenvalue equation in Eq. (B13), the eigenfrequencies and a set of orthogonal eigenvectors for the linearized Vlasov equation can be found numerically. If the two beams have the same lattice tune, the eigenfrequencies of Eq. (B13) are identical to that obtained in Ref. [4]. Fig. 8a is an example of eigenfrequencies of Eq. (B13) for the case of $\nu_{1,x} = \nu_{2,x}$ and $\xi_{1,x} = \xi_{2,x}$. It shows that in the symmetrical case of beam-beam interactions the coherent frequency (the first frequency line from the left of Fig. 8a) is separated from the rest of the eigenfrequencies that form a continuous band. The width of the band equals the incoherent beam-beam tune shift. The coherent

frequency can therefore be easily identified in this case. Note that the coherent beam-beam tune shift calculated from this coherent frequency is the same as that in Ref. [4]. The situation is more complicated when the two beams have very different lattice tunes. Fig. 8b plots the eigenfrequencies for the case of the HERA experiment. In this case, the eigenfrequencies are divided into two groups, one for each beam. For the e^+ beam, the eigenfrequencies form a continuous band that starts at the lattice tune of the e^+ beam and has a width of the incoherent beam-beam tune shift of the e^+ beam. Because there were two IPs in the HERA experiment, the incoherent beam-beam tune shift in Fig. 8b is $2\xi_{e,x} = 0.082$ for the e^+ beam. The characteristics of the eigenfrequencies for the p beam, in principle, is similar to that of the e^+ beam. Since the beam-beam parameter of the proton beam is very small ($\xi_{2,x} = \xi_{p,x} \sim 10^{-4}$), all the eigenfrequencies for the p beam degenerate into a single line (the first line from the left of Fig. 8b) that corresponds to the lattice tune of the p beam. In the case of very unsymmetrical beam-beam interactions, therefore, the coherent frequencies cannot be simply identified from the eigenfrequencies of the linearized Vlasov equation.

It should be noted that because of a very small beam-beam parameter of the p beam, during the HERA experiment no coherent beam-beam tune shift was observed on the proton beam. The matrix elements of $\mathbf{O}_2$ are very small as compared with the diagonal elements of $\mathbf{M}_1$ and $\mathbf{M}_2$. $\mathbf{O}_2$ can therefore be approximated as a zero matrix and the eigenfrequencies for the e^+ beam can be easily obtained from $\mathbf{M}_1 \vec{V}_1 = \nu \vec{V}_1$, where $\vec{V}_1 = (\bar{\psi}_{10}, \bar{\psi}_{11}, \dots, \bar{\psi}_{1l_p})^T$ is the sub vector space associated with the e^+ beam. Since $\mathbf{M}_1$ is diagonalized, solving the eigenfrequencies and a set of orthogonal eigenvectors of $\mathbf{M}_1$ is trivial. The eigenfrequencies and eigenvectors obtained from $\mathbf{M}_1$ were found to be the same as that of Eq. (B13) in the sub vector space associated with the e^+ beam in this case.

B.2. Initial-Value Problem for Coherent Frequencies

Let $(\nu_1, \dots, \nu_{l_p+1}, \nu_{l_p+2}, \dots, \nu_{2l_p+2})$ and $(\vec{V}^{(1)}, \dots, \vec{V}^{(l_p+1)}, \vec{V}^{(l_p+2)}, \dots, \vec{V}^{(2l_p+2)})$ be the eigenfrequencies and eigenvectors of the discretized and linearized Vlasov equation where $\mathbf{M}\vec{V}^{(n)} = \nu_n \vec{V}^{(n)}$. In the HERA experiment, the lattice tunes of the two beams are very different and,

therefore, the two eigenfrequency bands of Eq. (B13) are well separated (see Fig. 8b). In this case, $\{\nu_n\}$ are the eigenfrequencies for the e^+ beam when $n = 1, \dots, l_p + 1$ and the eigenfrequencies for the p beam when $n = l_p + 2, \dots, 2l_p + 2$. In the discretized action space, the perturbation of the beam distribution $\psi_1(I, \phi, \theta)$ and $\psi_2(I, \phi, \theta)$ can be represented as a vector,

$$\vec{\psi}(\theta) = (\psi_1(0, \phi, \theta), \psi_1(\Delta I, \phi, \theta), \dots, \psi_1(l_p \Delta I, \phi, \theta), \psi_2(0, \phi, \theta), \dots, \psi_2(l_p \Delta I, \phi, \theta))^T.$$

With the single-mode approximation, the general solution of $\psi_1(I, \phi, \theta)$ and $\psi_2(I, \phi, \theta)$ can then be obtained from a superposition of the eigenvectors of the linearized Vlasov equation,

$$\vec{\psi}(\theta) = \sum_{k=1}^{2l_p+2} C_k \vec{V}^{(k)} e^{i(\phi - \nu_k \theta)} \quad (B17)$$

where $\{C_k\}$ are constants and can be determined with an initial condition, $\psi_1(I, \phi, 0)$ and $\psi_2(I, \phi, 0)$. Since $|C_k|^2$ is the oscillation amplitude of the beam distributions with the frequency of ν_k , the diagram of $|C_k|^2$ v.s. ν_k corresponds to the frequency spectrum of the coherent oscillation. The two peaks in the $|C_k|^2$ - ν_k diagram, therefore, provides the coherent tunes when $\psi_1(I, \phi, 0) \rightarrow 0$ and $\psi_2(I, \phi, 0) \rightarrow 0$.

Consider a small kick that kicks beam 1 away from its equilibrium distribution $f_{1,0}(I, \phi)$, where $f_{1,0}(I, \phi)$ is known numerically from the beam-beam simulation. The initial perturbation of the beam distribution is

$$\psi_1(I, \phi, 0) = f_{1,0}(x + x_0, p_x) - f_{1,0}(x, p_x) = \sum_m g_m(I) e^{im\phi} \quad (B18)$$

and $\psi_2(I, \phi, 0) = 0$ where x_0 is the initial kick. With the single-mode approximation, $\psi_1(I, \phi, 0) \simeq g_1(I) e^{i\phi}$ and $\vec{\psi}(0) = \vec{g} e^{i\phi}$ where $\vec{g} = (g_1(0), g_1(\Delta I), \dots, g_1(l_p \Delta I), 0, \dots, 0)^T$. Note that the second half of the vector are all zero because beam 2 is not kicked. On the other hand, from Eq. (B17)

$$\vec{\psi}(0) = \left(\sum_{k=1}^{2l_p+2} C_k \vec{V}_k \right) e^{i\phi} = (\mathbf{V} \vec{C}) e^{i\phi} \quad (B19)$$

where $\mathbf{V}$ is a $(2l_p+2) \times (2l_p+2)$ matrix of which the i th column is $\vec{V}_i$ and $\vec{C} = (C_1, C_2, \dots, C_{2l_p+2})^T$. The coefficients $\{C_k\}$ can then be calculated from $\vec{C} = \mathbf{V}^{-1} \vec{g}$. It should be noted that the initial

kick on the beam distributions in Eq. (B18) can be in any direction in phase space since the coherent frequency is the frequency of an infinitesimal oscillation. For near-integrable systems considered in this study, the phase-space region in the vicinity of the origin is integrable and only consists of invariant circles (tori). It is therefore isotropic. The coherent frequencies calculated were indeed found to be independent of the direction of the initial kick.

Figure 8 plots the calculated $|C_k|^2\text{-}\nu_k$ diagrams for the symmetrical case of beam-beam interactions where $\nu_{1,x} = \nu_{2,x}$ and $\xi_{1,x} = \xi_{2,x}$ (Fig. 9a) and for the HERA experiment (Fig. 9b). In Fig. 9a, the peak with an arrow is the calculated coherent frequency that is the same as that in Fig. 8a. In Fig. 9b, the main peak indicates the calculated coherent frequency of the e^+ beam in the HERA experiment. The small peak in the lower right corner is the coherent frequency of the p beam. Figure 10 plots the calculated coherent beam-beam tune shift of the e^+ beam as a function of initial kick x_0 for the case of the HERA experiment. It shows that the calculated coherent beam-beam tune shift increases with the decrease of x_0 and converges as x_0 approaches zero. This amplitude dependence of the coherent frequency is consistent with the beam simulation. Since the coherent frequency is the frequency of an infinitesimal oscillation, the convergence of the calculated coherent frequency at $x_0 \rightarrow 0$ provides the wanted coherent frequency. As shown in our study (see Section IV), the linearized Vlasov equation with single-mode approximation is a valid approach for the coherent beam oscillation with symmetrical beam-beam interactions but not with unsymmetrical beam-beam interactions.

ACKNOWLEDGMENTS

This work is supported by the US Department of Energy under Grant No. DE-FG02-04ER41288. The authors would like to thank the Center for Advanced Scientific Computing at the University of Kansas for the use of the Supercomputer. The authors thank F. Willeke for many stimulating discussions. J. Shi would also like to thank DESY for the financial support for a trip to DESY at an early stage of our collaboration.

References

- [1] A.W. Chao and R.D. Ruth, Part. Accel. **16**, 201 (1985).
- [2] J. Shi and D. Yao, Phy. Rev. E**62**, 1258 (2000).
- [3] R.E. Meller and R.H. Siemann, IEEE Trans, NS-28, 2431 (1981).
- [4] K. Yokoya and H. Koiso, Part. Accel. **27**, 181 (1990), KEK Report 89-14 (1989), and the references herein.
- [5] K. Hirata, Nucl. Instr. Meth. **269**, 7 (1988).
- [6] H. Koiso, et al., Part. Accel. **27**, 83 (1990).
- [7] E. Keil, CERN Report, CERN-LEP-NOTE-294, (1981).
- [8] A. Hofmann, in *PROC. OF THE WORKSHOP ON BEAM-BEAM EFFECTS IN LARGE HADRON COLLIDERS*, edited by J. Poole and F. Zimmermann, CERN-SL-99-039 AP, (1999), p. 56.
- [9] A. Piwinski, IEEE Trans. NS-26, 4268 (1979).
- [10] R. Talman, CLNS-84/610, (1984).
- [11] Y. Alexahin, Part. Accel. **59**, 43 (1998).
- [12] M. Vogt, T. Sen, and J.A. Ellison, Phys. Rev. ST Accel. Beams **5**, 024401 (2002).
- [13] M.P. Zorano and F. Zimmermann, Phys. Rev. ST Accel. Beams **3**, 044401 (2000).
- [14] G.H. Hoffstaetter in *HERA Accelerator Studies 1999*, edited by G.H. Hoffstaetter, DESY HERA 00-02. (2000). p. 6.
- [15] L. Jin and J. Shi, Phy. Rev. E**69**, 036503 (2004).

- [16] G.H. Hoffstaetter in *HERA Accelerator Studies 2000*, edited by G.H. Hoffstaetter, DESY HERA 00-07. (2000). p. 5.
- [17] M. Sands, Stanford Linear Accelerator Center Report No. SLAC-121 (1970).
- [18] K. Hirata and F. Ruggiero, LEP Note 611, (1988).
- [19] B. S. Schmekel, G. H. Hoffstaetter, and J. T. Rogers, Phys. Rev. STAB**6**, 104403 (2003).

Table I. Some beam parameters used in HERA 2000 beam experiment. The horizontal size of the e^+ beam at the IPs is $\sigma_{e,x} = 283 \mu\text{m}$. The horizontal and vertical size of the p beam at the IPs are $\sigma_{p,x}=164 \mu\text{m}$ and $\sigma_{e,y}=39.9 \mu\text{m}$, respectively.

$\beta_{e,y}$ (m)	I_{e^+} (mA)	$\sigma_{e,y}(\mu\text{m})$	$\xi_{e,x}/\xi_{e,y}$	$\xi_{p,x}/\xi_{p,y}$ (10^{-4})
1.0	19	35.8	0.041/0.068	2.54/1.40
1.5	18	43.8	0.041/0.102	2.35/1.06
2.0	17	50.6	0.041/0.136	2.18/0.85
3.0	3.5	62.0	0.041/0.204	0.43/0.14
4.0	2.6	72.0	0.041/0.272	0.31/0.09

Table II. The coherent beam-beam tune shifts of the e^+ beam at $\beta_{e,y} = 4.0$ m ($\xi_{e,y} = 0.272$). “Experiment” and “Simulation” are the coherent tunes measured in the experiment and calculated in the beam simulation, respectively. “Rigid-Real” and “Rigid-Gaussian” are the coherent tunes calculated by using the rigid-beam model with a Gaussian distribution and with the distribution from the simulation, respectively. “Vlasov Eq.” is the coherent tunes calculated with the linearized Vlasov equation.

$\beta_{e,y} = 4.0$ m	$\nu_{e,x}$	$\Delta\nu_{e,x}/(2\xi_{e,x})$	$\nu_{e,y}$	$\Delta\nu_{e,y}/(2\xi_{e,y})$
Experiment	0.1600	0.110	0.2330	0.024
Simulation	0.1605	0.104	0.2331	0.024
Rigid-Real	0.1555	0.164	0.2194	0.049
Rigid-Gauss	0.1531	0.194	0.2040	0.077
Vlasov Eq.	0.1123	0.69		

Table III. The same as Table II, but for $\beta_{e,y} = 1.0$ m ($\xi_{e,y} = 0.068$).

$\beta_{e,y} = 1.0$ m	$\nu_{e,x}$	$\Delta\nu_{e,x}/(2\xi_{e,x})$	$\nu_{e,y}$	$\Delta\nu_{e,y}/(2\xi_{e,y})$
Simulation	0.1600	0.110	0.2172	0.212
Rigid-Real	0.1517	0.212	0.2074	0.284
Rigid-Gauss	0.1475	0.263	0.1996	0.341
Vlasov Eq.	0.121	0.58		

Table IV. The same as Table III, but with only one tenth of the p -bunch current used in the experiment ($\xi_{e,y} = 0.0068$).

$\beta_{e,y} = 1.0$ m	$\nu_{e,x}$	$\Delta\nu_{e,x}/(2\xi_{e,x})$	$\nu_{e,y}$	$\Delta\nu_{e,y}/(2\xi_{e,y})$
Simulation	0.1672	0.224	0.2414	0.337
Rigid-Real	0.1668	0.263	0.2406	0.396
Rigid-Gauss	0.1668	0.268	0.2406	0.399
Vlasov Eq.	0.165	0.53		

FIGURE CAPTIONS

Figure 1. Emittance of the e^+ beam as a function of $\xi_{e,y}$. ϵ_0 is the emittance without collision. Discrete points are from the experiment and continuous curves from the beam simulation. Circles and curve a is the vertical emittance. Crosses and curve b are the horizontal emittance. The two experimental data points at each $\xi_{e,y}$ where the measurement was performed correspond to the measurements at H1 and ZEUS.

Figure 2. The specific luminosity as a function of $\xi_{e,y}$. Circles are from the experiment and continuous curves from the beam simulation. The two experimental data points at each $\xi_{e,y}$ where the measurement was performed correspond to the measurements at H1 and ZEUS. Crosses are the luminosity calculated with the measured emittance in Fig. 1 assuming Gaussian beam distributions.

Figure 3. Evolution of the vertical emittance of the e^+ beam calculated with the beam simulation for the cases of HERA beam experiment at (a) $\beta_{e,y} = 1.0$ m; (b) $\beta_{e,y} = 1.5$ m; (c) $\beta_{e,y} = 2.0$ m; (d) $\beta_{e,y} = 3.0$ m; and (e) $\beta_{e,y} = 4.0$ m.

Figure 4. Power spectrum of the centroid motion of the e^+ beam in (a) horizontal and (b) vertical direction for the case of HERA beam experiment at $\xi_{e,y} = 0.272$ ($\beta_{e,y} = 4.0$ m). The beam centroid motion was calculated during the beam simulation. Note that in this case the incoherent beam-beam tune shifts for the two IPs are 0.082 and 0.544 in the horizontal and vertical plane, respectively.

Figure 5. The projection of the particle distribution of the e^+ beam in the vertical direction obtained by the beam simulation for the case of HERA beam experiment at $\xi_{e,y} = 0.272$ ($\beta_{e,y} = 4.0$ m). (a) The initial Gaussian distribution, (b) the distribution at the 5000th turn, and (c) the Gaussian distribution of which the standard deviation is the same as that of the beam distribution in (b).

Figure 6. The angle dependence of the particle distribution of the e^+ beam obtained by the beam simulation for the case of HERA beam experiment at $\xi_{e,y} = 0.272$ ($\beta_{e,y} = 4.0$ m). (a) The initial Gaussian distribution, (b) the ϕ_x -dependence and (c) the ϕ_y -dependence of the distribution

at the 5000th turn.

Figure 7. The ratio of the coherent and incoherent beam-beam tune shift as a function of the beam-beam parameter of the e^+ beam in the vertical plane calculated from (a) the beam-beam simulation and (b) the rigid-beam model with the Gaussian distribution.

Figure 8. Eigenfrequencies of Eq. (B13) for the case of (a) $\nu_{1,x} = \nu_{2,x}$, $\xi_{1,x} = \xi_{2,x}$ and (b) the HERA beam experiment. In (a), the dashed line marks the lattice tune that corresponds to the 0-mode. The single isolated line on the left is the coherent frequency that corresponds to a ratio of the coherent and incoherent beam-beam tune shift of 1.35. In (b), the dashed lines mark the lattice tunes of the e^+ beam (the left line) and p beam (the right line), respectively. The single isolated line on the right is the degenerated eigenfrequencies for the p beam and the band on the left is for the e^+ beam. The vertical axis has no physical meaning.

Figure 9. $|C_k|^2$ as a function of ν_k for the cases of Fig. 8. The arrows indicate the coherent frequencies.

Figure 10. Calculated coherent beam-beam tune shift by using the linearized Vlasov equation as a function of initial kick x_0 on the distribution of the e^+ beam [see Eq. (B15)] for the case of the HERA experiment at $\beta_{e,y} = 1.0$ m.

Figure 1




Figure 2




Figure 3




Figure 4a




Figure 4b




Figure 5




Figure 6




Figure 7




Figure 8a




Figure 8b




Figure 9a




Figure 9b




Figure 10

